

# A SHORT NOTE ON LIGAND FIELD THEORY OF THE MAGNETIC ANISOTROPY AND SUSCEPTIBILITY OF $\text{Fe}^{2+}$ TUTTON SALTS

B. BHATTACHARYYA\*

DEPT. OF MAGNETISM, INDIAN ASSOCIATION FOR THE CULTIVATION  
OF SCIENCE, CALCUTTA-32.

(Received May 26, 1964)

Investigations on the magnetic susceptibility and anisotropy of  $\text{Fe}(\text{NH}_4\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$  (Bose *et al.*, 1961) by Pryce's method shows a 20% reduction of the spin-orbit coupling owing to covalency effects. We have therefore, applied here the more reasonable and general method of molecular orbitals (Van-Vleck 1932, Stevens 1953, and Bose *et al.*, 1960) assuming that the covalency overlap is anisotropic.

The theoretical calculations of principal ionic susceptibilities of  $\text{Fe}^{2+}$  are made and compared with the experimental data on anisotropy and mean susceptibility for the salt  $\text{Fe}(\text{KSO}_4)_2 \cdot 6\text{H}_2\text{O}$  (Table I) which is isomorphous to the ammonium double salt previously investigated. It is assumed here as in the previous case that the ligand field has a small tetragonal component superposed on the predominant cubic part.

As no paramagnetic resonance data for  $\text{Fe}^{2+}$  Tutton salts is available, we compare the  $g$ -values of  $\text{Fe}(\text{KSO}_4)_2 \cdot 6\text{H}_2\text{O}$  with those of Tinkham's (1955) on  $\text{FeF}_2$  diluted with  $\text{ZnF}_2$ . Since the crystalline fields in  $\text{FeF}_2$  is truly orthorhombic the  $g$ -values are somewhat different in magnitude from that in Tutton salts but the order of magnitudes are comparable within the limits of approximations involved.

In the case of  $\text{Fe}(\text{NH}_4\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$ , the tetragonal field coefficient  $\Delta$  increases with temperature from a value of  $270 \text{ cm}^{-1}$  at  $20^\circ\text{K}$  to  $650^{-1} \text{ cm}$ . at  $300^\circ\text{K}$ . In the case of  $\text{Fe}(\text{KSO}_4)_2 \cdot 6\text{H}_2\text{O}$ ,  $\Delta$  changes from  $240^{-1} \text{ cm}$  at  $90^\circ\text{K}$  to  $640 \text{ cm}^{-1}$  at  $300^\circ\text{K}$ . Thus we see that the order of the anisotropic field and also their variations with temperature are almost the same in the two cases. The increase in  $\Delta$  with temperature is due to thermal expansion or relaxation effects in the salts. In the case of  $\text{Fe}(\text{NH}_4\text{SO}_4)_2 \cdot 6\text{H}_2\text{O}$  the spin-orbit coupling coefficient has been decreased by 20% from its free ion value of  $-103 \text{ cm}^{-1}$  (Owen 1955, Bose *et. al.*,

---

\*Lecturer, St. Xavier's College, Calcutta and Honorary Research Scholar, Indian Association for the Cultivation of Science.

1961). In the case of  $\text{Fe}(\text{KSO}_4)_2 \cdot 6\text{H}_2\text{O}$  we find the spin orbit coupling  $\zeta_{||} = -80 \text{ cm}^{-1}$ ,  $\zeta_{\perp} = -83 \text{ cm}^{-1}$  instead of free ion value of  $-103 \text{ cm}^{-1}$ . This indicates again almost the same overlap between the 3d  $\text{Fe}^{2+}$  and  $s$ - and  $p$ - $0^2$ -charge clouds, except that the overlap here has been taken as anisotropic.

The details of the work will be published shortly elsewhere.

TABLE I

$\text{Fe}(\text{KSO}_4)_2 \cdot 6\text{H}_2\text{O}$ .

$$k_{||} = 0.95, \quad \zeta_{||} = 0.80 \text{ cm}^{-1}$$

$$k_{\perp} = -80, \quad \zeta_{\perp} = -88 \text{ cm}^{-1}$$

Temp. °K	$\text{cm}^{-1}$	$p^2$	$p_{  }^2 - p_{\perp}^2$	$g$ -values
300	640	28.57 (28.58)	10.31 (10.34)	..
220	416	28.13 (28.08)	12.87 (12.70)	..
90	240	26.00 (26.27)	22.10 (22.01)	$g = 8.48$ $(8.97 \pm .02)$ $g = 6$ (6)

The values in the parenthesis in the 3rd and 4th column indicate the experimental results of Bose (1948). The  $g$ -values within parenthesis in the 5th column are Tinkham's (1955).

## ACKNOWLEDGMENT

The author is grateful to Prof. A. Bose, D.Sc., F.N.I. for suggesting the problem and guidance in the work. He is also grateful to his colleagues Sri R. Chatterjee, D.Phil., Sri U.S. Ghosh and Sri R. Rai of the department for all help and discussions in working out the theory and calculations of the parameters. Finally, he is very thankful to the authorities of the St. Xavier's College, Calcutta and Indian Association for the Cultivation of Science for permitting him to carry out the research project.

## REFERENCES

- Bose, A., 1948, *Ind. J. Physics*. **22**, 483.  
 Bose, A. Chakravarti A. S. and Chatterjee R, 1961, *proc. Roy. Soc. A* **261**, 207.  
 Bose, A. Chakravarti A. S. and Chatterjee R, 1960, *Proc. Roy. Soc. A*. **255**, 145.  
 Owen, J., 1955, *Proc. Roy. Soc. A*. **227**, 193.  
 Stevens K. W. H. 1953, *Proc. Roy. Soc. A*. **219**, 542.  
 Tinkham M. 1955, *Proc. Roy. Soc. A*. **236**, 535.  
 Van-Vleck J. H. 1932, *Phys. Rev.* **41**, 208.